



Research article

Breast cancer screening among Arabic women living in the State of Qatar: Awareness, knowledge, and participation in screening activities

Tam Truong Donnelly^{1,*}, Al-Hareth Al Khater², Salha Bujassoum Al-Bader², Mohamed Ghaith Al Kuwari³, Nabila Al-Meer², Mariam Malik³, Rajvir Singh², Sofia Chaudhry⁴, Roqia Ahmad Dorri⁴

¹Faculty of Medicine, The University of Calgary, P.O. Box 23133 Doha, Al Rayyan Campus, Al Forousiya Road, Qatar

²Hamad Medical Corporation

³Primary Health Care

⁴University of Calgary, Qatar

*Email: tdonnell@ucalgary.ca

ABSTRACT

Background: Breast cancer is the most common cancer among women in the State of Qatar, and the incidence rate is rising. Previous findings indicate women in Qatar are often diagnosed with breast cancer at advanced stages and their participation rates in screening activities are low.

Purpose: To investigate within the State of Qatar Arabic women's knowledge regarding breast cancer and breast cancer screening (BCS) methods and their participation rates in BCS. This paper reports on the results of a cross-sectional survey.

Methods: A quantitative, cross-sectional interview survey was conducted with 1,063 Arabic women (Qatari citizens and non-Qatari Arabic-speaking residents), 35 years of age or older, from March 2011 to July 2011.

Results: Of the 1,063 women interviewed (87.5% response rate), 90.7% were aware of breast cancer; 7.6% were assessed with having basic knowledge of BCS, 28.9% were aware of breast self-examination (BSE), 41.8% were aware of clinical breast exams (CBE), and 26.9% were aware of mammograms. Of the women interviewed, 13.8% performed BSE monthly, 31.3% had a CBE once a year or once every two years, and 26.9% of women 40 years of age or older had a mammogram once a year or once every two years. Participation rates in BCS activities were significantly related to awareness and knowledge of BCS, education levels, and receiving information about breast cancer, self-examination or mammography from any of a variety of sources, particularly physicians.

Conclusions: Study results demonstrate that despite the existent breast cancer screening recommendations, less than one-third of Arabic women living in Qatar participate in BCS activities. Public health campaigns encouraging more proactive roles for health care professionals regarding awareness and knowledge of breast cancer, BCS, and the benefits of early detection of breast cancer will help increase screening rates and reduce mortality rates among Arabic women living in the State of Qatar.

Keywords: Qatar, breast cancer screening, breast cancer awareness and knowledge, Arabic women breast cancer screening, Qatar females, breast self-examination, clinical breast examination, mammogram

<http://dx.doi.org/10.5339/avi.2012.2>

Published: 4 July 2012
© 2012 Donnelly et al., licensee Bloomsbury Qatar Foundation Journals. This is an open access article distributed under the terms of the Creative Commons Attribution License CC BY 3.0 which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Cite this article as: Donnelly TT, Al-Khater AH, Al-Bader SB, Al Kuwari MG, Al-Meer N, Malik M, Singh R, Chaudhry S, Dorri RA. Breast cancer screening among Arabic women living in the State of Qatar: Awareness, knowledge, and participation in screening activities, *Avicenna* 2012:2 <http://dx.doi.org/10.5339/avi.2012.2>

BACKGROUND

Worldwide, breast cancer is the most commonly diagnosed and one of the leading causes of cancer deaths [1,2]. In the Middle East region, the incidence of breast cancer has substantially increased in the last 24 years [3]. Women are not only being diagnosed with breast cancer in the later stages of the disease but also a higher proportion of younger women in their thirties and forties are clinically presenting with breast cancer [3–5]. According to the World Health Organization (WHO), Qatar experienced a 12.9/100,000 breast cancer mortality rate in 2008 [6]. This mortality rate is higher than other Arab peninsular countries, such as Saudi Arabia (10.4/100,000) and the United Arab Emirates (10.9/100,000), which also have lower incidence rates of breast cancer compared to the rest of the Middle East [6].

Qatar has experienced substantial development and lifestyle changes in the past few decades, which have been largely driven by oil and natural gas wealth [7]. Prevalence and patterns of cancer have been affected by rapid growth and subsequent environmental and social changes [3,7,8]. In 2007, 20% of cancer cases receiving treatment in one oncology hospital in Doha (Al Amal Hospital) were for breast cancer [9]. Data reviewed by Bener et al. from Qatar's 2006 National Cancer Disease Registry, and the WHO (2008) revealed that breast cancer is the leading cancer diagnosis, far greater than other most common cancers for Qatari women [6,7].

A note on terminology

We refer throughout the paper to 'Arabic' women. This term defines our study group as Arabic speakers, despite the fact that they are not all ethnically Arab. We use the term Arabic as a shorthand for 'Arabic-speaking' in this paper. See 'Study population' in the Methods section for further details.

Breast cancer incidence rates in the Middle East region

According to the International Agency for Cancer Research and GLOBOCAN 2008 estimates of age-standardized rates of breast cancer incidence and mortality, breast cancer incidence rates in the Arab Middle East are currently highest in Lebanon (55.4/100,000), followed by Bahrain (49.8/100,000) and Kuwait (47.7/100,000) [6]. In the Gulf Cooperation Council (GCC) countries,¹ breast cancer incidence rates are highest in Bahrain (49.8/100,000), followed by Kuwait (47.7/100,000) and Qatar (38.1/100,000). Compared to other Arab peninsular countries, such as Saudi Arabia (22.4/100,000) or Yemen (20.8/100,000), Qatar's incidence rate is high [6,10]. While breast cancer incidence rates in Western and European countries have stabilized or increased slightly in recent years [1,6,11], breast cancer incidence rates in Middle Eastern countries are increasing rapidly, and Qatar is no exception [7,8]. Bener et al. (2008) reviewed Qatar's National Cancer Disease Registry and found a 57.1% increase in total cancer cases in Qatar from 2002 to 2006 (360 cases compared to 170 cases between 1991 and 1996) [7].

Breast cancer screening

Mortality rates have decreased by 25 to 30% with early detection, improved quality of screening activities, and enhanced treatment [11,12]. Breast cancer detected in the early stages has a higher chance of responding successfully to treatment [12,13]. Arabic women currently face a significant risk of high mortality rate from breast cancer due to frequent diagnosis in the advanced stages of the disease [7,8,14,15].

Early detection and screening activities of breast cancer include breast self examination (BSE), clinical breast examination (CBE), and mammography. Mammography can reduce mortality rates in women aged 40 to 74 by 25% [11,16]. The WHO (2011) reports mammography to be the most successful way of detecting breast cancer amongst women older than 50 [1].

Previously in Qatar, the breast cancer screening (BCS) recommendation was for a monthly BSE with a yearly CBE for women aged above 35 years and a mammogram every two years for women aged 40–69 years, unless otherwise advised by physicians [17,18]. However, BCS recommendations are currently being revised in Qatar to reflect Qatar's health care context and to meet the need of cancer care for younger women. The guidelines now recommend a monthly BSE starting at age 20, a yearly

¹GCC countries include Saudi Arabia, the United Arab Emirates (UAE), Qatar, Bahrain, Oman and Kuwait.

CBE for women aged 35 and above, and an annual mammogram for women aged 40–69 years, unless otherwise advised by physicians.

Although data related to breast cancer research in the Middle East is scarce [7,19], previous studies in the region have found low BCS participation rates amongst women [8,13,20–22]. In a more recent study of 1,200 Qatari national women between the ages of 30–55 years, it was discovered that despite an adequate knowledge of breast cancer, participation rates in breast cancer screening activities were low; BSE 24.9%, CBE 23.3% and mammography 22.5% [13,20]. Of note, women taking part in these activities were mainly young and of higher socioeconomic status. These rates are very low when compared to women's breast cancer screening rates in countries such as Canada, Denmark, Finland, Sweden and the Netherlands, where 70% to 85% women aged 50–69 years have had a mammogram in the previous three years [23]. National screening programs vary; therefore, benchmarking remains a challenge. However, low breast cancer screening participation rates in Qatar should raise concern among health care providers and policy makers alike.

Barriers to Arabic women's breast cancer screening

Factors that are associated with women's compliance or lack thereof in BCS activities must be investigated to create an effective intervention strategy. Studies have also shown that physician's recommendation acts as an enabler, while not receiving a recommendation can act as a barrier [8]. Previous studies have revealed that additional barriers to mammogram practice include fear of cancer, fear of finding out one has cancer, the notion that there is no cure, perceived benefit, time, cost, fear of gossip, fear that breast examination and mammogram could be painful, husband or other male family members objecting to breast examination, preference for a female health professional, accessibility of the health care system, perceived effectiveness and embarrassment [8,14,20–22,24,25].

In some conservative Arab areas, barriers to BCS can also include travel restrictions for women, some of whom may not drive or travel alone without a male due to laws or customs [20,22]. Coupled with male relatives who may not be supportive or who object to such examinations, women are vulnerable to the risk of discovering breast cancer at later stages if they are not comfortable raising issues of breast lumps and breast exams. Consequently, health promotion messages must also promote male encouragement of their female relatives' decision to seek a mammogram [19].

Studies have shown that knowledge, attitudes, beliefs and practices related to breast cancer screening practices are influenced by social and cultural frameworks [8,14,20–22,24,25]. Therefore, it is essential to examine knowledge, attitudes, beliefs, practices, barriers and enablers for specific populations before designing and implementing an intervention strategy [21,25–27]. Accordingly, contextual factors that influence breast cancer screening practices amongst Arabic women living in the State of Qatar must be investigated in order to design a socio-culturally appropriate intervention strategy that addresses the threat of breast cancer in Qatar.

The multiple intervention framework

Because breast cancer screening is complex in nature and has many interrelated determinants, promoting breast cancer screening behaviors requires a comprehensive set of intervention strategies targeting individual behavior changes, collective capacity and action, and supportive environments. The Canadian Ottawa Charter for Health Promotion confirms that health is influenced by contextual factors including those that are social, economic, political, and cultural [28]. Thus, developing a successful health promotion program necessitates more complex, multi-sector interventions that facilitate individual behavioral changes and address those contextual factors that influence health [28,29]. See Fig. 1.

The design of multiple intervention programs for community health involves the deployment of multiple strategies via multiple channels to deal with the health problems. These strategies are aimed at individuals, groups, communities, organizations and policies. The goal of this research study is to develop, implement, and sustain an intervention program that raises awareness of breast cancer and increases women's participation in breast cancer screening activities. To achieve this goal, a multiple intervention conceptual framework was adopted for the design of our research program [29]. A **three-phase research program** was planned. The aim of Phase I was to seek understanding of breast health issues as experienced by Arabic women in Qatar. The aim of **Phase II** is to identify and




Figure 1. Multiple intervention framework [29].

implement strategies that would assist and encourage women to participate in breast cancer screening activities. The aim of **Phase III** is to evaluate the feasibility and sustainability of such strategies in order to increase participation in breast cancer screening activities over the long term [30–32].

Phase I currently consists of two studies: a quantitative cross-sectional interview survey and an exploratory, qualitative study using individual in-depth interviews conducted with Arabic women, Arabic men and health care providers in Qatar. Based on the results of the quantitative cross-sectional survey, the primary outcomes of interest measured for this paper include (a) awareness of and participation rates in breast cancer and BCS of Arabic women living in Qatar, and (b) the relationship between Arabic women’s awareness, knowledge of breast cancer and its screening, select socio-demographic factors, and their participation in BCS (mammogram, clinical breast examination, breast self examination).

As part of phase I of our research program, a quantitative, descriptive cross-sectional community-based interview survey with Arabic women living in Qatar was conducted from March 2011 to July 2011 to investigate women’s participation rates in breast cancer screening activities, their knowledge about breast cancer and its screening methods, barriers and facilitators to participation, and their view of the current breast cancer screening programs in the State of Qatar. Data was collected to answer the following *research questions*:

1. What are the participation rates of Arabic women in BSE, CBE, and mammogram?
2. To what extent are Arabic women’s awareness and knowledge of breast cancer (BC) and its screening, as well as other selected factors, associated with BCS behaviors?

Ethics approval for this research study was obtained from the Hamad Medical Corporation Research Committee and the University of Calgary’s Conjoint Health Research Ethics Board. Consent to participate in this study was obtained from each participant. Prior to conducting an interview, each participant was given an explanation of the study and her rights according to standard interview guidelines. Participants were assured that all information would remain confidential, and interview questionnaires were stripped of identifying information to preserve confidentiality.

METHODS

Study population

According to 2010 census data in Qatar [33], there are approximately 72,240 women who are 35 years of age and older living in Doha, Al Wakrah, and Al Khor/Al Thakhira, of which there are 60,937 women 35 years and older in Doha; 7,909 in Al Wakrah; and 3,394 in Al Khor/Al Thakira. Based on a 95% confidence level, the study’s sample size was calculated using Cochran’s formula for sample size [34]. See Fig. 2.

A convenience sample of 1,215 self-identified Arabic women (Qatari citizens and non-Qatari Arabic-speaking residents) who met the study’s inclusion criteria was approached and invited to

City	Population of women 35 years and over	Sample size using a margin of error of 3.5%	Sample size using a margin of error of 5%
Doha	60,937	640	315
Al Wakrah	7,909	83	41
Al Khor	3,394	36	18
Total	72,240	759	374

Figure 2. Study sample calculation based on Cochran's formula for sample size.

participate in the survey, 1,063 of whom agreed to participate in a structured, face-to-face interview (87.5% response rate). Study participant inclusion criteria also included being aged 35 years or older, recruitment from one of seven different health care settings in Qatar, and having lived in Qatar for at least 10 years. Women who did not meet the mentioned inclusion criteria were excluded from the study. To ensure that information gathered reflected the experiences of women living in both urban and semi-urban/rural areas, participants were recruited from urban hospital settings and community health clinics in Doha, Al Wakrah (South of Qatar), and Al Khor (North of Qatar). Other main sites in Qatar are currently under Qatar Petroleum and Qatar Gas jurisdiction. Not only is accessibility to these sites difficult, but Qatar Petroleum and Qatar Gas are responsible for the provision of health care to its workers, with the majority of these workers being male.

Questionnaire and data collection method

Face-to-face interviews using a structured, quantitative survey questionnaire were used for data collection. Interviews were conducted in Arabic by seven female interviewers and lasted an average of 30 minutes. These interviewers were nurses who were fluent in both Arabic and English, had experience working in the health care system in Qatar, and were trained by the lead principal investigator of the project (TD). Five of these research staff had a Bachelor of Science degree in Nursing and two were in their final year of obtaining the degree. To ensure diversity of participants and represent the general female Arabic population, study participants were interviewed during different days of the week and different times of the day.

Survey items had been developed specifically for this study and were further refined based on the responses and suggestions of experts in the field. The structured questions were based upon the literature and the results of previous studies [35–40]. With permission from researchers who conducted similar surveys with women of different ethno-cultural backgrounds—Dr. Stephen McPhee [41–43] and Dr. Juliane Cheek [44]—items from their surveys of breast cancer screening with Vietnamese women living in the U.S and Australia, respectively, were also incorporated. These items have been tested and proven valid and reliable. Back-translation of the survey questionnaire into Arabic and English was carried out to ensure lexical equivalence. The survey instrument was field tested with 30 Arabic women (35 years of age and older) at the various research sites by a female interviewer fluent in both Arabic and English prior to conducting the full-scale study; these results were not included in the full-scale study, and were meant to further clarify and refine questionnaire items, develop community connections and coalition, and assess the feasibility of the full-scale survey.

Statistical analysis

Descriptive statistics in the form of mean, standard deviations for interval variables and frequency, with percentages for categorical variables, were performed. To see associations between dependent and categorical variables, chi-square tests were performed. Co-linearity for all covariates significant at univariate analysis was tested before using them in the multivariate logistic regression analysis.² Multivariate logistic regression was used to assess the effects of selected socio-demographic factors, awareness and knowledge of BCS, health care provider information, and sources of information about breast cancer and mammogram in predicting the binary dependent variables (e.g. clinical breast

²The CBE Awareness variable was kept in the multivariate analysis, despite its high colinearity with CBE practice.

examination, breast self examination, mammogram). A P value of 0.05 (two tailed) was considered as a statistically significant level. Data analysis was done using statistical program SPSS version 19.

RESULTS

Participants' profile: overall demographics

The 1,063 participants who completed the interview were between the ages of 35–82 ($M = 44.9$, $SD = 8.4$) years old; 78.9% were married, 98.2% were Muslims, 84.6% had children (49.2% had five or more children), 88.7% resided in urban areas, 33.3% attended university, 30.9% reported their husband attended university, 34.1% were employed and 58.8% were full-time homemakers.

According to 2010 Qatar Census results, Qatari citizens represent 24.4% of the population in Qatar; Qatari female citizens represent 36.7% of the female population living in Qatar [33]. Overall, 52.1% of the women interviewed in this study were Qatari citizens, 10.9% were citizens of other Arab peninsular/GCC countries (Saudi Arabia, UAE, Kuwait, Oman, Bahrain, Yemen), 16% were citizens of Levant countries (Syria, Lebanon, Palestine, Jordan), 10.1% were citizens of North African countries (Egypt, Libya, Tunisia, Algeria, Morocco), and 10.9% were citizens of other countries in the greater Middle East (Sudan, Iraq, Iran, Somalia, Mauritania, Pakistan). All participants completed the interview in Arabic and resided in Qatar for at least 10 years, with the majority living in Qatar between 30–49 years ($M = 34.8$, $SD = 14.6$).

Using the U.S. Bureau of Labor Statistics Standard Occupational Classification system to categorize participants' occupations, more than two-thirds of employed participants (69.1%) were categorized as working in the "Management, Business, Science or Arts" field [45]. Forty-three (4%) of the women interviewed reported that they had been diagnosed with breast cancer (Table 1).

Awareness of breast cancer and breast cancer screening activities

In this study, the term "awareness" is used to indicate that a participant has a preliminary understanding of breast cancer and its screening, whereas the term "basic knowledge" indicates that a participant has a higher level of understanding and accurate knowledge of breast cancer and its screening. For example, a participant was assessed as having breast cancer awareness if she had heard of breast cancer and received some information about breast cancer from at least one source (Table 2). However, a participant was further assessed as having basic knowledge of breast cancer screening only if she knew how to examine her own breasts, and knew the most recent national screening recommendations for mammography, breast self-examination and clinical breast examination (Table 3). It is important to note that the current study's indicators for assessing participants' awareness and basic knowledge were different (as indicated in Tables 2 and 3) than in other studies conducted in the region [13,22].

To qualify as being aware, participants had to meet the criteria for each category listed in Table 2. The present study found that while 90.7% of participants had awareness of breast cancer, only 28.9% had awareness of BSE, 41.8% of CBE, and 26.4% of mammography.

Basic knowledge of breast cancer screening activities

Upon assessing higher level knowledge of screening recommendations, only 7.6% of participants were assessed with having basic knowledge of breast cancer screening activities (Table 3).

Breast cancer screening participation rates

For the current study, appropriate practice of breast cancer screening activities was defined according to the most recent national screening guidelines and local health center recommendations; breast self examination (BSE) must have been performed by the participant monthly, clinical breast examination (CBE) must have been performed by a health care professional (most often a doctor or a nurse) within the last 1–2 years, and participants 40 years of age or older must have had a mammogram within the last 1–2 years.

The current study found 13.9% of participants reported performing a monthly BSE, 31.3% had a CBE within the last 1 to 2 years, and 26.9% of participants 40 years of age or older had a mammogram within the last 1–2 years. Over 50% of the participants interviewed reported they had never participated in BCS activities (Table 4). Almost half of survey participants reported that they plan to have a CBE or mammogram in the next 12 months; 49.4% of the participants said they plan on having a breast exam by a health care professional, and 47.8% of the participants ≥ 40 years old

Table 1. Selected socio-demographic data (N = 1,063)

Variable	N (%)
Age (<i>M</i> = 44.9, <i>SD</i> = 8.4)	
35–39 years	365 (34.4)
40–49 years	399 (37.6)
50–59 years	220 (20.7)
60+ years	77 (7.2)
Nationality	
Qatari (Qatari citizen)	554 (52.1%) ^a
Non-Qatari (Qatari resident)	509 (47.9%)
Religion	
Muslim	1044 (98.2%)
Christian	19 (1.8%)
Marital status	
Single/never married	83 (7.8%)
Married	839 (78.9%)
Other (separated, divorced, widowed)	141 (13.3%)
Education level of participant^b	
Never went to school	146 (13.7%)
Primary school	213 (20%)
High school	287 (27%)
Trade school	62 (5.8%)
University	354 (33.3%)
Employment status of participant^b	
Employed	362 (34.1%)
Management, Business, Science, Arts	250 (69.1%)
Service Occupations	34 (9.4%)
Sales and Office Occupations	64 (17.5%)
Construction, Maintenance Occupations	1 (0.3%)
Production, Transportation Occupations	4 (1.1%)
Full time homemaker	625 (58.8%)
Unemployed	75 (7.1%)
Education level of husband	
Never went to school	112 (10.5%)
Primary school	164 (15.4%)
High school	233 (21.9%)
Trade school	59 (5.6%)
University	328 (30.9%)
Not applicable/other	167 (15.7%)
Living area	
Urban	943 (88.7%)
Semi-urban	120 (11.3%)
Years in Qatar (<i>M</i> = 34.8, <i>SD</i> = 14.6)	
10–29 years	332 (31.2)
30–49 years	551 (51.8)
50+ years	180 (16.9)
Has participant ever had breast cancer?	
Yes	43 (4%)
No	1020 (96%)

^a According to 2010 Qatar Census results, Qatari citizens represent 24.4% of the entire population in Qatar. Qatari female citizens represent 36.7% of the entire female population in Qatar [33].

^b 1 participant did not answer these questions.

said they plan on having a mammogram. Only 1.5% of women participants reported that their husbands or male family members are not supportive of them having breast examinations by a health care professional, and 1.0% reported their husbands or other family members' objection was the reason they were not planning on having a mammogram.

Relationship between awareness, knowledge, educational levels and participation rates

Bivariate analyses using chi-square testing were also used to assess relationships between awareness, knowledge, educational levels and participation rates in BCS activities (Table 5). The current study found awareness and basic knowledge of BCS amongst participants were significantly related to participation rates in breast cancer screening activities. Participants who had appropriately practiced BSE were more likely to have breast cancer awareness, $\chi^2(1, N = 1062) = 8.74$, $p = 0.003$; BSE awareness, $\chi^2(1, N = 1063) = 199.53$, $p = 0.001$; CBE awareness,

Table 2. Awareness of breast cancer and breast cancer screening (N = 1063).

Awareness of Breast Cancer Participant must have correctly answered both questions:	Correct n (%)	Incorrect n (%)
Have you ever heard of breast cancer? [Yes]	993 (93.4%)	70 (6.6%)
Have you ever had any information about breast cancer from: [participant must say “Yes” to at least one]:		
Family member or friend	713 (67.1%)	350 (32.9%)
Newspapers/magazines	692 (65.2%)	371 (34.8%)
Television	781 (73.5%)	282 (26.5%)
Pamphlet	505 (47.6%)	558 (52.4%)
Doctor	305 (28.7%)	758 (71.3%)
Nurse	185 (17.4%)	878 (82.6%)
Health educator	176 (16.6%)	887 (83.4%)
Number (%) of participants who correctly answered both questions	964 (90.7%)	
Awareness of Breast Self Examination Participant must have correctly answered both questions:	Correct n (%)	Incorrect n (%)
Have you ever heard of women performing breast self-examination at home? [Yes]	436 (41.0%)	627 (59.0%)
Do you know how to examine your own breast? [Yes]	469 (44.1%)	594 (55.9%)
Number (%) of participants who correctly answered both questions	307 (28.9%)	
Awareness of Clinical Breast Examination Participant must have correctly answered both questions:	Correct n (%)	Incorrect n (%)
Have you ever heard of an exam where a doctor or a nurse examines a woman’s breast to feel for a small lump that could be an early sign of breast cancer? [Yes]	753 (70.8%)	310 (29.2%)
Have you ever had such a breast examination by a doctor or a nurse? [Yes]	451 (42.4%)	612 (57.6%)
Number (%) of participants who correctly answered both questions	444 (41.8%)	
Awareness of Mammography Participant must have correctly answered both questions:	Correct n (%)	Incorrect n (%)
At what age does the Qatar screening program suggest that women should start having mammograms? [40–50]	387 (36.4%)	676 (63.6%)
How often does the screening program suggest that women should have a mammogram? [once a year, once every two years]	509 (47.9%)	554 (52.1%)
Number (%) of participants who correctly answered both questions	281 (26.4%)	

Table 3. Basic knowledge of breast cancer screening (N = 1, 063).

Participant must have correctly answered all 6 questions:	Correct n (%)	Incorrect n (%)
At what age does the Qatar screening program suggest that women should start having mammograms? [40–50]	387 (36.4%)	676 (63.6%)
How often does the screening program suggest that women should have a mammogram? [once a year, once every two years]	509 (47.9%)	554 (52.1%)
Have you ever heard of an exam where a doctor or a nurse examines a woman’s breast to feel for a small lump that could be an early sign of breast cancer? [Yes]	753 (70.8%)	310 (29.2%)
Have you ever had such a breast examination by a doctor or a nurse? [Yes]	451 (42.4%)	612 (57.6%)
Have you ever heard of women performing breast self-examination at home? [Yes]	436 (41.0%)	627 (59.0%)
Do you know how to examine your own breast? [Yes]	469 (44.1%)	594 (55.9%)
Number (%) of participants who correctly answered all 6 questions	81 (7.6%)	

$\chi^2(1, N = 1063) = 91.28, p = 0.001$; mammogram awareness, $\chi^2(1, N = 1062) = 11.44, p = 0.001$; and basic knowledge of BCS, $\chi^2(1, N = 1063) = 79.63, p = 0.001$. Similar trends were found for appropriate practices of CBE and mammogram (with p values ≤ 0.05).

Higher education levels of participants were also significantly related with appropriate practice of BCS activities [BSE, $\chi^2(1, N = 1062) = 31.14, p = 0.001$; CBE, $\chi^2(1, N = 1062) = 16.16, p = 0.003$;

Table 4. Breast cancer screening participation rates.

Participant practiced BCS activities	At appropriate time ^a	At some point	Never practiced
		n (%)	n (%)
Breast Self Examination (BSE) (N = 1063) ^b	148 (13.9%)	294 (27.7%)	621 (58.4%)
Clinical Breast Examination (CBE) (N = 1063)	333 (31.3%)	118 (11.1%)	612 (57.6%)
Mammogram (only women aged ≥ 40) (N = 696) ^c	187 (26.9%)	87 (12.5%)	421 (60.6%)

^a Although breast cancer screening guidelines are being revised to reflect Qatar's health context and to meet the need of cancer care for women, appropriate times in this study indicate BSE Monthly; CBE for women aged ≥ 35 within the last 1–2 years; mammogram for women aged ≥ 40 within the last 1–2 years.

^b 5 participants did not answer this question.

^c 1 participant did not answer this question. Therefore, calculations related to mammogram practice are based on $n = 695$.

Mammogram, $\chi^2(1, N = 695) = 18.36, p = 0.001$. This was also true for their husbands' education levels for participants who were ever married [BSE, $\chi^2(1, N = 896) = 38.65, p = 0.001$; CBE, $\chi^2(1, N = 896) = 17.15, p = 0.002$; Mammogram, $\chi^2(1, N = 604) = 15.34, p = 0.004$].

Demographic factors such as age, nationality, marital status, residential information, and employment status were also analyzed (Table 6). Marital status was significantly related to CBE practice, $\chi^2(2, N = 1063) = 20.51, p = 0.001$. Employed participants were significantly more likely to perform BSE monthly than unemployed participants, [$\chi^2(2, N = 1062) = 16.07, p = 0.001$]. While age did not significantly relate to appropriate mammogram practice, the 40–49 year old age group was significantly more likely to have performed a monthly BSE [$\chi^2(3, N = 1061) = 17.57, p = 0.001$] or had a CBE within the last 1–2 years [$\chi^2(3, N = 1061) = 24.92, p = 0.001$] than other age groups. Similarly, participants who had been living in Qatar for 50 years or more were significantly less likely to perform monthly BSE, [$\chi^2(2, N = 1063) = 17.26, p = 0.001$]; however, length of residence did not affect CBE or mammogram practice. It is more likely that older Qatari women are less aware of or know how to perform breast self examination. Living area (urban versus semi-urban areas) was also not significantly related to participation in BCS (Table 6).

Multivariate analysis of factors associated with practice of breast cancer screening

Table 7 represents the multivariate logistic regression analysis of selected independent factors that may be used as indicators to predict participation in breast cancer screening activities. Variables with a small number of cases were removed from the multivariate regression analysis.

Participants who were aware of BSE or CBE had between two to seven times the odds of performing a BSE appropriately than those without awareness (OR = 6.40; 95% CI = 3.97–10.33; $p < 0.001$ and OR = 2.78; 95% CI = 1.63–4.73; $p < 0.001$, respectively). Having awareness of CBE was the strongest predictor for appropriate CBE practice (OR = 185.56; 95% CI = 81.50–422.50; $p < 0.001$). Participants whose doctor talked to them about breast cancer had two to three times the odds of practicing BSE (OR = 2.08; 95% CI = 1.05–4.14; $p = 0.037$) or having CBE (OR = 3.52; 95% CI = 1.63–7.61; $p = 0.001$) than those whose doctors did not. Participants had increased odds of having a BSE or CBE if they received information about breast cancer from a nurse (CBE: OR = 2.72; 95% CI = 1.29–5.75; $p = 0.009$), newspaper/magazine (BSE: OR = 2.10; 95% CI = 1.05–4.18; $p = 0.035$), or received information about mammography from a doctor (CBE: OR = 1.74; 95% CI = 1.02–2.95; $p = 0.041$), or television/radio (CBE: OR = 2.03; 95% CI = 1.03–3.99; $p = 0.041$) than those who did not.

Although having mammogram awareness was not a predictor for having a mammogram, participants with CBE awareness were more likely to have had a mammogram at appropriate/recommended times (OR = 6.51; 95% CI = 3.65–11.63; $p < 0.001$). The strongest predictor for having a mammogram was receiving information about a mammogram from a doctor (OR = 10.10; 95% CI = 5.85–17.44; $p < 0.001$). Participants also had at least twice the odds of having a mammogram if their doctor talked to them about breast cancer (OR = 3.15; 95% CI = 1.35–7.40; $p = 0.008$), or if they received information about mammography from a pamphlet (OR = 2.26; 95% CI = 1.08–4.70; $p = 0.030$).

Table 5. Awareness, knowledge, education and participation in breast cancer screening activities.

Variable	Appropriate practice of BSE		Appropriate practice of CBE		Participation in Mammogram	
	Yes (%) N=148	No (%) N=915	Yes (%) N=333	No (%) N=730	Yes (%) N=187	No (%) N=508
Awareness of breast cancer	144 (97.3)	820 (89.6)	324 (97.3)	640 (87.7)	180 (96.3)	441 (86.8)
	$\chi^2(1, N = 1062) = 8.74, p = 0.003$		$\chi^2(1, N = 1062) = 26.81, p = 0.001$		$\chi^2(1, N = 695) = 12.82, p = 0.001$	
Awareness of BSE	115 (77.7)	192 (21.0)	163 (48.9)	144 (19.7)	80 (42.8)	116 (22.8)
	$\chi^2(1, N = 1063) = 199.53, p = 0.001$		$\chi^2(1, N = 1063) = 95.08, p = 0.001$		$\chi^2(1, N = 695) = 26.86, p = 0.001$	
Awareness of CBE	115 (77.7)	329 (36.0)	326 (97.9)	118 (16.2)	164 (87.7)	164 (32.3)
	$\chi^2(1, N = 1063) = 91.28, p = 0.001$		$\chi^2(1, N = 1063) = 628.10, p = 0.001$		$\chi^2(1, N = 695) = 168.44, p = 0.001$	
Awareness of mammogram	56 (37.8)	225 (24.6)	129 (38.7)	152 (20.8)	82 (44.1)	91 (17.9)
	$\chi^2(1, N = 1062) = 11.44, p = 0.001$		$\chi^2(1, N = 1062) = 38.14, p = 0.001$		$\chi^2(1, N = 694) = 49.84, p = 0.001$	
Basic knowledge of BCS	38 (25.7)	43 (4.7)	66 (19.8)	15 (2.1)	33 (17.6)	25 (4.9)
	$\chi^2(1, N = 1063) = 79.63, p = 0.001$		$\chi^2(1, N = 1063) = 102.53, p = 0.001$		$\chi^2(1, N = 695) = 28.94, p = 0.001$	
Education participant						
Never went to school	8 (5.4)	138 (15.1)	29 (8.7)	117 (16.0)	20 (10.7)	112 (22.0)
Primary school	16 (10.8)	197 (21.6)	60 (18.0)	153 (21.0)	40 (21.4)	122 (24.0)
High School	39 (26.4)	248 (27.1)	90 (27.0)	197 (27.0)	46 (24.6)	127 (25.0)
Trade school	11 (7.4)	51 (5.6)	23 (6.9)	39 (5.3)	15 (8.0)	29 (5.7)
University	74 (50.0)	280 (30.6)	131 (39.3)	223 (30.6)	66 (35.3)	118 (23.2)
	$\chi^2(4, N = 1062) = 31.14, p = 0.001$		$\chi^2(4, N = 1062) = 16.16, p = 0.003$		$\chi^2(4, N = 695) = 18.36, p = 0.001$	
Education husband						
Never went to school	3 (2.4)	109 (14.1)	23 (7.8)	89 (14.8)	16 (9.8)	87 (19.8)
Primary school	9 (7.2)	155 (20.1)	42 (14.2)	122 (20.3)	25 (15.2)	89 (20.2)
High School	31 (24.8)	202 (26.2)	85 (28.7)	148 (24.7)	43 (26.2)	106 (24.1)
Trade school	13 (10.4)	46 (6.0)	23 (7.8)	36 (6.0)	10 (6.1)	29 (6.6)
University	69 (55.2)	259 (33.6)	123 (41.6)	205 (34.2)	70 (42.7)	129 (29.3)
	$\chi^2(4, N = 896) = 38.65, p = 0.001$		$\chi^2(4, N = 896) = 17.15, p = 0.002$		$\chi^2(4, N = 604) = 15.34, p = 0.004$	

DISCUSSION AND RECOMMENDATIONS

The results of the current study demonstrates that the majority of Arabic women in Qatar are aware of breast cancer (90.7%), however, there are much lower levels of awareness and knowledge of breast cancer screening activities, and only 7.6% of the women interviewed were assessed with having basic knowledge of the most recent screening guidelines in Qatar for BSE, CBE and mammograms.

Previous studies conducted throughout the Middle East [5,8,13–15,20–22] have also found that BCS participation rates are low. Bener et al. (2009) in a study of 1,200 Qatari women (aged 30–55) found 24.9% of those interviewed practice BSE, 23.3% had CBE, and 22.5% had mammography [13]. The current study assessed participation rates in breast cancer screening activities according to the BCS recommended guidelines in Qatar. As expected, current findings

Table 6. Selected demographic factors and participation in breast cancer screening activities.

Variable	Appropriate practice of BSE		Appropriate practice of CBE		Participation in Mammogram	
	Yes (%) N=148	No (%) N=915	Yes (%) N=333	No (%) N=730	Yes (%) N=187	No (%) N=508
Age Groups						
35–39	47 (32.2)	318 (34.8)	88 (26.5)	277 (38.0)	-	-
40–49	75 (51.4)	324 (35.4)	150 (45.2)	249 (34.2)	110 (58.8)	289 (56.9)
50–59	20 (13.7)	200 (21.9)	80 (24.1)	140 (19.2)	64 (34.2)	155 (30.5)
60+	4 (2.7)	73 (8.0)	14 (4.2)	63 (8.6)	13 (7.0)	64 (12.6)
	$\chi^2(3, N = 1061) = 17.57, p = 0.001$		$\chi^2(3, N = 1061) = 24.92, p = 0.001$		$\chi^2(2, N = 695) = 4.62, p = 0.099$	
Nationality						
Qatari Citizens	69 (46.6)	485 (53.0)	163 (48.9)	391 (53.6)	98 (52.4)	280 (55.1)
Qatari Residents	79 (53.4)	430 (47.0)	170 (51.1)	339 (46.4)	89 (47.6)	228 (44.9)
	$\chi^2(1, N = 1063) = 2.08, p = 0.149$		$\chi^2(1, N = 1063) = 1.95, p = 0.163$		$\chi^2(1, N = 695) = 0.41, p = 0.524$	
Marital status						
Single	14 (9.5)	69 (7.5)	11 (3.3)	72 (9.9)	4 (2.1)	25 (4.9)
Married	119 (80.4)	720 (78.7)	289 (86.8)	550 (75.3)	158 (84.5)	392 (77.2)
Others	15 (10.1)	126 (13.8)	33 (9.9)	108 (14.8)	25 (13.4)	91 (17.9)
	$\chi^2(2, N = 1063) = 1.92, p = 0.383$		$\chi^2(2, N = 1063) = 20.51, p = 0.001$		$\chi^2(2, N = 695) = 5.15, p = 0.076$	
Living area						
Urban	137 (92.6)	806 (88.1)	298 (89.5)	645 (88.4)	171 (91.4)	440(86.6)
Semi-Urban	11 (7.4)	109 (11.9)	35 (10.5)	85 (11.6)	16 (8.6)	68 (13.4)
	$\chi^2(1, N = 1063) = 2.55, p = 0.110$		$\chi^2(1, N = 1063) = 0.29, p = 0.588$		$\chi^2(1, N = 695) = 3.00, p = 0.083$	
Years in Qatar						
10–29	65 (43.9)	267 (29.2)	111 (33.3)	221 (30.3)	59 (31.6)	134 (26.4)
30–49	71 (48.0)	480 (52.5)	169 (50.8)	382 (52.3)	84 (44.9)	238 (46.9)
50+	12 (8.1)	168 (18.4)	53 (15.9)	127 (17.4)	44 (23.5)	136 (26.8)
	$\chi^2(2, N = 1063) = 17.26, p = 0.001$		$\chi^2(2, N = 1063) = 1.09, p = 0.579$		$\chi^2(2, N = 695) = 1.98, p = 0.371$	
Employment status						
Employed	71 (48.0)	291 (31.8)	124 (37.2)	238 (32.6)	57 (30.5)	131(25.8)
Homemaker	72 (48.6)	553 (60.5)	182 (54.7)	443 (60.8)	112 (59.9)	333 (65.7)
Unemployed	5 (3.4)	70 (7.7)	27 (8.1)	48 (6.6)	18 (9.6)	43 (8.5)
	$\chi^2(2, N = 1062) = 16.07, p = 0.001$		$\chi^2(2, N = 1062) = 3.62, p = 0.164$		$\chi^2(2, N = 694) = 2.00, p = 0.367$	

indicate that participation rates in BCS activities among Arabic women living in Qatar were also low; only 13.9% of participants performed a monthly breast self examination (BSE), 31.3% had a clinical breast examination (CBE) within the last 1 to 2 years, and 26.9% of participants 40 years of age or older had a mammogram done within the last 1 to 2 years. Compared to European guidelines set to >70% acceptance levels of non-opportunistic breast cancer screening programs, participation rates in Qatar and other Gulf Cooperation Countries (GCC) are very low [10,46]. This study's results also show that there are some women who have, at some point, performed BSE (27.7%), had CBE (11.1%) and mammogram (12.5%). Together with findings that indicate almost half of participants said that they plan to have a CBE (49.4%) or mammogram (47.8%, >40 years old) within the next 12 months, these are encouraging findings. It may be that with encouragement and more awareness of national guidelines, more Qatari women may practice BCS according to the recommended guidelines.

Table 7. Multivariate logistic regression analysis of selected factors associated with practice of breast cancer screening.

Variables	Category	OR	95% CI	P value
Appropriate practice of BSE				
BSE awareness	Yes	6.40	3.97–10.33	<0.001
CBE awareness	Yes	2.78	1.63–4.73	<0.001
Doctor talked to participant about breast cancer	Yes	2.08	1.05–4.14	0.037
Participant received information about breast cancer from a newspaper/magazine	Yes	2.10	1.05–4.18	0.035
Appropriate practice of CBE				
CBE awareness	Yes	185.56	81.50–422.50	<0.001
Doctor talked to participant about breast cancer	Yes	3.52	1.63–7.61	0.001
Participant received information about breast cancer from a nurse	Yes	2.72	1.29–5.75	0.009
Participant received information about mammography from a doctor	Yes	1.74	1.02–2.95	0.041
Participant received information about mammography from television/radio	Yes	2.03	1.03–3.99	0.041
Appropriate practice of mammogram				
CBE awareness	Yes	6.51	3.65–11.63	<0.001
Doctor talked to participant about breast cancer	Yes	3.15	1.35–7.40	0.008
Participant received information on mammogram from a doctor	Yes	10.10	5.85–17.44	<0.001
Participant received information on mammogram from a pamphlet	Yes	2.26	1.08–4.70	0.030

It is important to note that the authors of this paper are aware of the controversial issue regarding the effectiveness and usefulness of breast self-examination for women living in other societies [47]. Given that the population of interest is women who come from different ethno-cultural backgrounds throughout the Middle East and who live in an area with notably low participation rates in CBE and mammography, *not* recommending BSE to Arabic women warrants very careful consideration of such a decision's impact on early detection of breast cancer in Qatar.

Both bivariate analyses and multivariate regression analyses showed nationality was not a significant predictor for BCS practice. We suggest this result is due to the fact that the Qatari government provides subsidized or free health care services—including mammograms at recommended times—in modern facilities for all of its residents, which reduces the burden of cost as a barrier to having a mammogram. Qatar is a multi-national country (the current study's sample population represented over 20 different nationalities); it is encouraging that nationality does not affect participation rates in BCS activities. However, since fewer Qatari women citizens reported that their doctors talked to them about breast cancer (19.3%) than other Arabic women living in Qatar (29.9%), those behind public education campaigns and health care providers should find ways to talk to and encourage Qatari women about breast cancer awareness and recommended screening activities. Previous studies also indicate that physicians are less likely to share information or recommend mammography to women different than themselves in nationality, age, gender or social class [8,27,48].

Although living area (urban versus semi-urban) was not significantly related to participation in BCS activities, living in semi-urban areas might limit the ability of women age 40 and older to access mammography screening clinics in urban areas because they have to travel longer distances to a mammogram clinic. Thus, a mobile mammogram clinic that provides service closer to women living in semi-urban areas may increase women's mammogram participation rates [49].

Participants with higher levels of education were more likely to participate in BCS activities. A recent study in Saudi Arabia (2011) indicates similar results; women with at least university level education are more likely to participate in BSE and CBE than women with lower educational levels [50]. It is important to note that not only did participation in breast cancer screening activities increase with higher education levels of women participants, but they also increased with higher education levels of their husbands. To our knowledge, no other studies have examined the relationship between the husband's level of education and the wife's breast cancer screening

knowledge and activities in the Middle East. The current study found husbands' higher education levels are significantly related to women having knowledge of BCS and to higher participation rates in breast cancer screening activities. It may be that men who have higher education levels are more aware of breast cancer, its screening, and the benefits of early detection of breast cancer; thus they may give more support and encouragement to their wives to participate in BCS activities. As men play an important role in Arabic women's lives, their interest and participation in breast cancer screening could be an effective enabler [19,49]. Thus, Arabic men should be made aware of breast cancer and its screening, and their role in providing support and encouragement to their female family members in BCS activities. It is a very encouraging finding from our study that only between 1.0 and 1.5% of participants perceived that their husbands or other family members object to breast examinations.

The current study also indicates receiving information about breast cancer and BCS from any source significantly increased participation in screening activities. Multivariate regression analyses indicate receiving information from a doctor about breast cancer was a predictor for performing breast self-examination, having a clinical breast examination or a mammogram at recommended times. Despite this, doctors do not often talk about breast cancer or actively encourage their female patients to participate in breast cancer screening activities; only 24.4% of the women interviewed reported that their doctors had ever talked to them about breast cancer. Women whose doctors talked to them about mammography were more likely to have a mammogram according to recommended guidelines. Previous studies have also shown physician recommendations are important determinants of mammography [27,50–52]. This indicates the critical role doctors have in raising awareness in women about breast cancer and breast cancer screening services.

Policies related to screening guidelines and patient education should stress increased physician-patient time to discuss breast cancer, screening recommendations, the benefits of early detection, and the negative effects of not participating in appropriate BCS activities. As current findings suggest, women are more likely to participate in BCS activities if they received information from even one source about breast cancer or mammography, and one of the strongest predictors of participating in BCS was receiving information from their doctors. Primary care physicians often build a rapport with their patients, and are therefore in an ideal position to raise awareness among women during routine patient visits [50]. Physicians should be encouraged to incorporate breast cancer screening recommendations into their daily routine and practice with their female patients.

The current study demonstrates that not only doctors, but also other health care providers such as nurses and health educators play a significant role in breast health care in Qatar. Women who received information about breast cancer from a nurse had twice the odds of having a CBE at recommended times than women who did not receive information from a nurse (OR = 2.72; 95% CI = 1.29–5.75; $p = 0.009$). Yet, only 17.4% of the women interviewed in this study received information about breast cancer from a nurse, 16.6% from a health educator, and 11.9% and 9% received information about mammography from nurses or health educators, respectively. Health care providers were found to be important sources of knowledge about breast cancer screening for Arabic women in other studies [50–56]. Several studies have also found that nurses and health care workers do not have sufficient knowledge of breast cancer risk factors and screening methods [57–60], and their lack of knowledge limits encouragement of patients to engage in screening activities [60]. Allied health professionals should therefore be trained and empowered to play a bigger and more proactive part in breast cancer intervention strategies in Qatar and neighboring countries. A 2011 study conducted in Qatar found that even pharmacists are willing to be involved with educating their patients about breast cancer, despite their current low involvement [61].

A Train-the-Trainer (TTT) program that focused on teaching breast cancer screening and prevention to nurses from 20 different countries indicates health care providers such as nurses and allied health professionals can help address knowledge gaps [62]. A similar study in Turkey found knowledge and participation rates of Turkish women increased following a BSE educational intervention program [63]. Although promising, rates are still low and indicate having knowledge may not be enough. Previous studies have found that even among Arab women with sufficient knowledge, participation in breast cancer screening activities remained low [13,22,53,57,64,65]. More research on socio-cultural barriers is needed for development of culturally appropriate interventions.

Given these findings—that there are low levels of awareness and knowledge of breast cancer screening and that women who are aware of BCS recommendations are more likely to practice BCS activities according to recommended guidelines—it is imperative that public educational campaigns

be implemented in Qatar to promote breast health and early detection of breast cancer. Breast cancer awareness campaigns are effective educational tools; they have been shown to increase motivation in women and improve physician initiative for early detection [19]. Raising awareness about breast cancer and its screening among women in Qatar through a variety of sources is also important when considering an intervention program. This study found women who received information about breast cancer or mammograms from any of the sources asked (family/friend, newspaper/magazine, television/radio, pamphlets, doctor, nurse, health educator) were more likely to participate in BCS. Although only 1.1% of the women interviewed listed the internet as their source of information on breast cancer, it is also important to utilize the internet to raise awareness about breast cancer and its screening among younger women. Sending reminders in the form of a letter or SMS message might be effective in increasing attendance at Well-Women clinics for breast examination and mammogram uptake [14,15]; 58.7% of women interviewed in the current study said that they would have a mammogram if they received a letter reminding them from a doctor or other health care provider.

CONCLUSION

While breast cancer incidence rates are rising among women in the Middle East [3,7,8], current findings on the strongest predictors for participation in breast cancer screening activities among Arabic women in Qatar are encouraging. An overwhelming majority (94%) of the women interviewed reported that they would make an appointment for a mammogram if their doctor recommended it, and a doctor recommendation is one of the strongest predictors of screening practices. It is therefore critical to raise awareness among physicians and other health care professionals such as nurses, health educators, or pharmacists of the effectiveness of their explanations and demonstrations to patients about breast cancer and its screening. It is also important to raise awareness of breast cancer and early detection amongst women through various forms of media to increase participation in early screening programs and reduce breast cancer morbidity and mortality for Arab women living in Qatar.

Due to convenience sampling, the current study has limited ability to generalize survey results and potential bias. To avoid bias to some extent, randomly-selected times were chosen for the face-to-face interviews, and attempts were made to approach all potential respondents in an interview location. This method has been used successfully in other studies to recruit respondents in hard-to-reach communities [66,67]. In addition, data was collected from self-reports, and may be subject to inaccuracy or social-desirability response bias.

Despite these limitations, the benefits of this study's findings are potentially far-reaching and include: raising awareness levels of breast cancer and its screening activities; developing a culturally appropriate, socially-acceptable, and effective intervention program; and ultimately, decreasing both morbidity and mortality from breast cancer in Qatar. Linking research findings to policy making, service delivery and practice is an important element of this study. As several investigators of this study are key health care providers and health care policy-makers who are committed to providing services that meet the health care needs of women in Qatar, we anticipate that the results of this research will impact health care policy as well as health care provision to women in Qatar. Findings may also be applicable to other female populations in the Middle East and benefit women of similar ethnic and cultural backgrounds worldwide.

COMPETING INTERESTS

The authors declare they have no competing interests.

FUNDING SOURCES

Qatar National Research Fund, National Priority Research Program (NPRP).
Project ID: NPRP 09-261-3-059

AUTHORS' CONTRIBUTIONS

Donnelly, T.; Al-Khater, A.; Al-Kuwari, M.; Singh, R.; Bujassoum Al-Bader, S.; Al-Meer, N.; Malik, M.; Chaudhry, S.; Dorri, R.

(i) DT: Contributed to the conception and design of the study and the acquisition, analysis and interpretation of the data, drafted the manuscript and gave final approval of the manuscript version submitted for publication.

(ii) AKA: Contributed to the conception and design of the study and the acquisition of the data, revised the manuscript and gave final approval of the manuscript version submitted for publication.

(iii) AKM: Contributed to the conception and design of the study and the acquisition of the data, review the manuscript critically for content and gave final approval of the manuscript version submitted for publication.

(iv) SR: Contributed to the conception and design of the study and the acquisition, analysis and interpretation of the data, revised the manuscript and gave final approval of the manuscript version submitted for publication.

(v) BAS: Contributed to the conception and design of the study and the acquisition of the data, revised the manuscript and gave final approval of the manuscript version submitted for publication.

(vi) AMN: Contributed to the conception and design of the study and the acquisition of the data, revised the manuscript and gave final approval of the manuscript version submitted for publication.

(vii) MM: Contributed to the conception and design of the study and the acquisition of the data, review the manuscript critically for content and gave final approval of the manuscript version submitted for publication.

(viii) CS: Contributed to the analysis and interpretation of the data, drafted the manuscript and gave final approval of the manuscript version submitted for publication.

(ix) DR: Contributed actively to the acquisition of data, organized workshop training for interviewers, liaison with each data collection site, translated material from English to Arabic.

ACKNOWLEDGEMENTS

This publication was made possible by a grant from Qatar National Research Fund under its National Priority Research Program (NPRP 09-261-3-059). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of Qatar National Research Fund. We are grateful to all the women who participated in this research, as well as to the Qatar National Research Fund, which provided us with funding to conduct this study. We also give special thanks to staff at the Hamad Medical Corporation (Hamad General Hospital, Women Hospital) and Qatar Primary Health Care, Community Health Clinics who helped us to recruit the research participants, all our research assistants (Roqaiya Ahmad Dorri, Aisha Al-Ali, Aisha Al-Khayren, Asma Albulushi, Asma Rehman, Fadi Al-Massri, Khadra Yassin, Salah Hmaid, Yasser Sami, Zeinab Idris, Noora Rashid Al Enazi, Nahridda Nazir Khiyal Meer), and our former project manager, Floor Christie de Jong.

References

- [1] WHO, Breast cancer: Prevention and control, 2011 [<http://www.who.int/cancer/detection/breastcancer/en/index.html>].
- [2] Jemal A, Bray F, Center MM, Ferlay J, Ward E and Forman D. Global cancer statistics. *CA Cancer J Clin.* 2011 Mar-Apr;61:2, 69–90.
- [3] Tarabeia J, Baron-Epel O, Barchana M, Liphshitz I, Ifrah A, Fishler Y and Green MS. A comparison of trends in incidence and mortality rates of breast cancer, incidence to mortality ratio and stage at diagnosis between Arab and Jewish women in Israel, 1979–2002. *Eur J Cancer Prev.* 2007 Feb;16:1, 36–42.
- [4] Al-Saad S, Al-Shinnawi H and Mahmood Shamsi N. Risk factors of breast cancer in Bahrain. *Bahrain Medical Bulletin.* 2009;31:2, 1–11.
- [5] Miller AB. Screening for breast cancer in the Eastern Mediterranean Region. *East Mediterr Health J.* 2010 Oct;16:10, 1022–1024.
- [6] IARC International Agency for Research Cancer, WHO, World cancer report 2008 and Global cancer statistics. [<http://www.iarc.fr/en/publications/pdfs-online/wcr/2008/index.php>] or [<http://globocan.iarc.fr/factsheet.asp>].
- [7] Bener A, Ayub H, Kakil R and Ibrahim W. Patterns of cancer incidence among the population of Qatar: a worldwide comparative study. *Asian Pac J Cancer Prev.* 2008 Jan–Mar;9:1, 19–24.
- [8] Azaiza F and Cohen M. Health beliefs and rates of breast cancer screening among Arab women. *J Womens Health (Larchmt).* 2006 June;15:5, 520–530.
- [9] Chanpong GF. Qatar World Health Survey 2006 Overview. Department of Public Health National Health Authority, Qatar (2008).
- [10] Al-Hamdan N, Ravichandran K, Al-Sayyad J, Al-Lawati J, Khazal Z, Al-Khateeb F, Abdulwahab A and Al-Asfour A. Incidence of cancer in Gulf Cooperation Council countries, 1998–2001. *East Mediterr Health J.* 2009 May–June;15:3, 600–611.
- [11] Mai V, Sullivan T and Chiarelli AM. Breast Cancer Screening Program in Canada: Successes and Challenges. *Salud Publica Mex* 2009. 51
- [12] Tabár L, Vitak B, Chen HH, Duffy SW, Yen MF, Chiang CF, Krusemo UB, Tot T and Smith RA. The Swedish Two-County Trial twenty years later. Updated mortality results and new insights from long-term follow-up. *Radiol Clin North Am.* 2000 July;38:4, 625–651.
- [13] Bener A, El Ayoubi HR, Moore MA, Basha B, Joseph S and Chouchane L. Do we need to maximise the breast cancer screening awareness? Experience with an endogamous society with high fertility. *Asian Pac J Cancer Prev.* 2009 Oct–Dec;10:4, 599–604.

- [14] Soskolne V, Marie S and Manor O. Beliefs, recommendations and intentions are important explanatory factors of mammography screening behavior among Muslim Arab women in Israel. *Health Educ Res.* 2007 Oct;22:5, 665–676.
- [15] Baron-Epel O, Friedman N and Lernau O. Reducing disparities in mammography-use in a multicultural population in Israel. *Int J Equity Health.* 2009 May;19:8, 19.
- [16] U S National Cancer Institute Mammograms [<http://www.cancer.gov/cancertopics/factsheet/detection/mammograms>].
- [17] Qatar National Cancer Society [<http://www.qncs.org.qa/breast.htm>].
- [18] Hamad Medical Corporation. [www.hmc.org.qa; <http://www.hmc.org.qa/hmcnewsite/news.aspx?id=606>].
- [19] El Saghir NS, Khalil MK, Eid T, El Kinge AR, Charafeddine M, Geara F, Seoud M and Shamseddine AI. Trends in epidemiology and management of breast cancer in developing Arab countries: a literature and registry analysis. *Int J Surg.* 2007 Aug;5:4, 225–233.
- [20] Bener A, Honein G, Carter AO, Da'ar Z, Miller C and Dunn EV. The determinants of breast cancer screening behavior: a focus group study of women in the United Arab Emirates. *Oncol Nurs Forum.* 2002 Oct;29:9, E91-8.
- [21] Azaiza F and Cohen M. Between traditional and modern perceptions of breast and cervical cancer screenings: a qualitative study of Arab women in Israel. *Psychooncology.* 2008 Jan;17:1, 34–41.
- [22] Bener A, Alwash R, Miller CJ, Denic S and Dunn EV. Knowledge, attitudes, and practices related to breast cancer screening: a survey of Arabic women. *J Cancer Educ.* 2001 Winter;16:4, 215–220.
- [23] WHO World Health Statistics. 2008 [http://www.who.int/whosis/whostat/EN_WHS08_Full.pdf].
- [24] Sadler GR, Dhanjal SK, Shah NB, Shah RB, Ko C, Anghel M and Harshburger R. Asian Indian women: knowledge, attitudes and behaviors toward breast cancer early detection. *Public Health Nurs.* 2001 Sep–Oct;18:5, 357–363.
- [25] Baron-Epel O. Attitudes and beliefs associated with mammography in a multiethnic population in Israel. *Health Educ Behav.* 2010 Apr;37:2, 227–242.
- [26] Bigby J. Global understanding of barriers to breast cancer screening. *J Womens Health (Larchmt).* 2006 June;15:5, 484.
- [27] Shirazi M, Champeau D and Talebi A. Predictors of breast cancer screening among immigrant Iranian women in California. *J Womens Health (Larchmt).* 2006 June;15:5, 485–506.
- [28] Epp J. Ottawa: Health and Welfare Canada; 1986. *Achieving Health for All: A Framework for Health Promotion.* [<http://www.frcentre.net/library/AchievingHealthForAll.pdf>].
- [29] Edwards N, Mill J and Kothari AR. Multiple intervention research programs in community health. *Can J Nurs Res.* 2004 Mar;36:1, 40–54.
- [30] Sampson R and Morenoff J. Public health and safety in context: Lessons from community-level theory on social capital. *Promoting Health: Intervention Strategies from Social and Behavioral Research.* Eds. Smedley BD and Syme SL 2000; National Academy Press, Washington. 366–389.
- [31] Edwards N. Ratcheting up research efforts for systems change. *Nurs Leadersh (Tor Ont).* 2003;16:2, 27–31.
- [32] Mill J and Edwards N. Incorporating culture and gender into multiple intervention programs. *Can J Nurs Res.* 2003 June;35:2, 9–14.
- [33] Qatar Census. Final Results of Census 2010 [<http://www.qsa.gov.qa/QatarCensus/Populations.aspx>].
- [34] Cochran WB. *Sampling Techniques.* Third ed, Wiley 1977.
- [35] Donnelly TT, McKellin W, Hislop G and Long B. Socioeconomic influences on Vietnamese-Canadian women's breast and cervical cancer prevention practices: a social determinant's perspective. *Soc Work Public Health.* 2009 Sep–Oct;24:5, 454–76.
- [36] Donnelly TT. Challenges in providing breast and cervical cancer screening services to Vietnamese Canadian women: the healthcare providers' perspective. *Nurs Inq.* 2008 June;15:2, 158–168.
- [37] Donnelly TT and McKellin W. Vietnamese Canadian women's breast cancer and cervical cancer screening: the influence of gendered roles and expectations. *Anthropology & Aging Quarterly.* 2008;29:3, 76–88.
- [38] Donnelly TT and McKellin W. Keeping healthy! Whose responsibility is it anyway? Vietnamese Canadian women and their healthcare providers' perspectives. *Nurs Inq.* 2007 Mar;14:1, 2–12.
- [39] Donnelly TT. The health-care practices of Vietnamese-Canadian women: cultural influences on breast and cervical cancer screening. *Can J Nurs Res.* 2006 Mar;38:1, 82–101.
- [40] Donnelly TT. Vietnamese women living in Canada: Contextual factors affecting Vietnamese women's breast cancer and cervical cancer screening practices. Doctoral Dissertation. 2004. The University of British Columbia, Canada: Faculty of Interdisciplinary Studies.
- [41] McPhee SJ, Stewart S, Brock KC, Bird JA, Jenkins CN and Pham GQ. Factors associated with breast and cervical cancer screening practices among Vietnamese American women. *Cancer Detect Prev.* 1997;21:6, 510–521.
- [42] McPhee SJ and Nguyen TT. Cancer, cancer risk factors, and community-based cancer control trials in Vietnamese Americans. *Asian Am Pac Isl J Health.* 2000 Winter;8:1, 18–31.
- [43] McPhee SJ, Nguyen TT, Shema SJ, Nguyen B, Somkin C, Vo P and Pasick R. Validation of recall of breast and cervical cancer screening by women in an ethnically diverse population. *Prev Med.* 2002 Nov;35:5, 463–473.
- [44] Cheek J, Fuller J, Gilchrist S, Maddock A and Ballantyne A. Vietnamese women and Pap smears: issues in promotion. *Aust N Z J Public Health.* 1999 Feb;23:1, 72–76.
- [45] U S Bureau of Labor Statistics. 2010 SOC User Guide [http://www.bls.gov/soc/soc_2010_user_guide.pdf].
- [46] Perry N, Broeders M, de Wolf C, Törnberg S, Holland R and von Karsa L. European guidelines for quality assurance in breast cancer screening and diagnosis. Fourth edition—summary document. *Ann Oncol.* 2008 Apr;19:4, 614–622.
- [47] Thomas DB, Gao DL, Ray RM, Wang WW, Allison CJ, Chen FL, Porter P, Hu YW, Zhao GL, Pan LD, Li W, Wu C, Coriaty Z, Evans I, Lin MG, Stalsberg H and Self SG. Randomized trial of breast self-examination in Shanghai: final results. *J Natl Cancer Inst.* 19;2:94, 1445–1457.
- [48] Keinan-Boker L. Performance of breast cancer early detection in Arab women in Israel—room for improvement. *J Womens Health (Larchmt).* 2006 Jun;15:5, 542–5.

- [49] Remennick L. The challenge of early breast cancer detection among immigrant and minority women in multicultural societies. *Breast J.* 2006 Jan–Feb;12:Suppl 1, S103-10.
- [50] Ravichandran K, Al-Hamdan NA and Mohamed G. Knowledge, attitude, and behavior among Saudis toward cancer preventive practice. *J Family Community Med.* 2011 Sep;18:3, 135–142.
- [51] Cohen M and Azaiza F. Early breast cancer detection practices, health beliefs, and cancer worries in Jewish and Arab women. *Prev Med.* 2005 Nov–Dec;41:5–6, 852–858.
- [52] Alam AA. Knowledge of breast cancer and its risk and protective factors among women in Riyadh. *Ann Saudi Med.* 2006 Jul–Aug;26:4, 272–7.
- [53] Montazeri A, Vahdaninia M, Harirchi I, Harirchi AM, Sajadian A, Khaleghi F, Ebrahimi M, Haghighat S and Jarvandi S. Breast cancer in Iran: need for greater women awareness of warning signs and effective screening methods. *Asia Pac Fam Med.* 2008 Dec 20;7:1, 6.
- [54] Al-Qattan M, Al Saleh K, Al-Musallam S and Masoud G. Knowledge and factors affecting breast self examination. *Kuwait Medical Journal.* 2007;40:2.
- [55] Ahmed BA. Awareness and practice of breast cancer and breast-self examination among university students in Yemen. *Asian Pac J Cancer Prev.* 2010;11:1, 101–105.
- [56] Dandash KF and Al-Mohaimed A. Knowledge, attitudes, and practices surrounding breast cancer and screening in female teachers of Buraidah, Saudi Arabia. *Int J Health Sci (Qassim).* 2007 Jan;1:1, 61–71.
- [57] Alkhasawneh IM, Akhu-Zaheya LM and Suleiman SM. Jordanian nurses' knowledge and practice of breast self-examination. *J Adv Nurs.* 2009 Feb;65:2, 412–416.
- [58] Haji-Mahmoodi M, Montazeri A, Jarvandi S, Ebrahimi M, Haghighat S and Harirchi I. Breast self-examination: knowledge, attitudes, and practices among female health care workers in Tehran. *Iran Breast J.* 2002 Jul–Aug;8:4, 222–225.
- [59] Jaradeen N. Breast cancer risk-factors and breast self examination practice among Jordanian women. *Bahrain Medical Bulletin.* 2010;32:1,
- [60] Madanat H and Merrill RM. Breast cancer risk-factor and screening awareness among women nurses and teachers in Amman, Jordan. *Cancer Nurs.* 2002 Aug;25:4, 276–282.
- [61] El Hajj MS and Hamid Y. Breast cancer health promotion in Qatar: a survey of community pharmacists' interests and needs. *Int J Clin Pharm.* 2011 Feb;33:1, 70–79.
- [62] Dow Meneses K and Yarbrow CH. Cultural perspectives of international breast health and breast cancer education. *J Nurs Scholarsh.* 2007;39:2, 105–12.
- [63] Gürsoy AA, Hindistan S, Nural N, Kahrman I, Yilmaz F, Yigitbas C, Erdöl H, Bulut HK, Calik KY and Mumcu HK. Comparison of three educational interventions on breast self-examination knowledge and health beliefs. *Asian Pac J Cancer Prev.* 2009;10:5, 765–772.
- [64] Dündar PE, Ozmen D, Oztürk B, Haspolat G, Akyildiz F, Coban S and Cakiroglu G. The knowledge and attitudes of breast self-examination and mammography in a group of women in a rural area in western Turkey. *BMC Cancer.* 2006 Feb 24;6:43.
- [65] Taha H, Halabi Y, Berggren V, Jaouni S, Nyström L, Al-Qutob R and Wahlström R. Educational intervention to improve breast health knowledge among women in Jordan. *Asian Pac J Cancer Prev.* 2010;11:5, 1167–1173.
- [66] CDC AIDS Community Demonstration Projects Research Group. The CDC AIDS Community Demonstration Projects: A multi-site community-level intervention to promote HIV risk reduction. *American Journal of Public Health.* 1999;89:3, 336–345.
- [67] Watters JK and Biernacki P. Targeted sampling: Options and considerations for the study of hidden populations. *Social Problems.* 1989;36:416–430.